



Received: 2016/12/20, Revised: 2017/03/08,  
 Accepted: 2017/03/18, Published: 2017/06/30

©2017 Yun-Seok Kang et al.; Licensee Journal of Exercise Nutrition and Biochemistry. This is an open access article distributed under the terms of the creative commons attribution license (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*Corresponding author: Jae-Cheol Kim  
 Department of Sports Science, Chonbuk National University, Jeonju, Republic of Korea.

Tel : +82-10-7104-2863 Email : [kjc@jbn.ac.kr](mailto:kjc@jbn.ac.kr)

©2017 The Korean Society for Exercise Nutrition

## Effects of swimming exercise on high-fat diet-induced low bone mineral density and trabecular bone microstructure in rats

Yun-Seok Kang<sup>1</sup>/ Sang-Hyun Kim<sup>1</sup>/ Jae-Cheol Kim<sup>1\*</sup>

1. Department of Sports Science, Chonbuk National University, Jeonju, Republic of Korea.

### INTRODUCTION

Bones are active living tissues that are sensitive not only to external stimuli such as exercise but also to metabolic changes within the body, and actively undergo metabolism<sup>26</sup>. During bone metabolism, resorption by osteoclasts is balanced by formation by osteoblasts, to repair microdamage and replace old tissues with new ones, and to maintain bone health<sup>23</sup>.

Bone metabolism is regulated by the differentiation of marrow mesenchymal stem cells, which can differentiate into osteoblasts and adipocytes<sup>6</sup>. From childhood to early adulthood, which is the period of increase in bone mass, the differentiation of marrow mesenchymal stem cells into osteoblasts increases owing to the effects of growth factors; however, during the period of a decrease in bone mass as a result of aging, differentiation into osteoblasts decreases, while differentiation into adipocytes increases; this phenomenon inhibits normal bone metabolism<sup>18</sup>. Osteopenia and osteoporosis are characterized by reduced bone mineral density, and can result from such metabolic changes in the bone<sup>17</sup>. Moreover, excessive accumulation of fat, as observed in obesity, induces the expression of various inflammatory cytokines, and promotes the differentiation of marrow mesenchymal stem cells and hematopoietic stem cells into osteoclasts to increase bone resorption and decrease bone mineral density, ultimately increasing the risk of osteoporosis and fractures<sup>20</sup>. Sawant *et al.* reported lower bone mineral density of the tibia and femur in rat models of high-fat diet-induced obesity than in normal rats<sup>29</sup>. This result indicates that high-fat diet-induced obesity has negative effects on bone mineral density.

Studies have been performed to identify methods to inhibit reduction in bone mineral density and stabilize or improve the existing levels. Among these methods, exercise is reported to not only provide appropriate stimuli for enhancing bone health but also increase the expression of growth factors, while reducing that of inflammatory factors by inducing weight loss, thereby positively affecting bone metabolism and bone health<sup>13, 26</sup>. Weight-bearing exercises such as running on a treadmill have most commonly been used in research<sup>12, 21, 27</sup>. However, several recent studies suggest that swimming can also effectively improve bone mineral density<sup>7, 14</sup>. Ju *et al.* reported increased femoral density and trabecular size after 12 weeks of

**[Purpose]** This study aimed to investigate the effect of swimming exercise on high-fat diet-induced low bone mineral density (BMD) and trabecular bone microstructure in rats.

**[Methods]** Eight-week-old male Sprague–Dawley (SD) rats were divided into a normal diet group (n=9) and a high-fat diet group (n=15). Three rats in each group were sacrificed after 8 weeks of high-fat diet to evaluate the association between high-fat diet and bone health. The other 18 rats were reasigned to 3 groups (normal diet control, NC; high-fat diet control, HC; high-fat diet + Exercise, HEx) for up to another 8 weeks. Rats in the exercise group were trained for a swimming exercise program (1 h/day, 5 times/week for 8 weeks). All rats were sacrificed 24 h after the last bout of exercise to analyze the BMD and trabecular bone microstructure in the femur and tibia, using micro-computed tomography.

**[Results]** First, the effect of high-fat diet on bone health was examined. It was observed that BMD, percent bone volume (BV/TV), and trabecular number (Tb.N) of the femur and tibia were lower in rats in the high-fat diet group than in those in the normal diet group (p < .05). In addition, BMD, BV/TV, and Tb.N of the femur and tibia were significantly increased in rats that underwent the 8-week swimming exercise program, compared to the corresponding values in rats in the HC group (p < .05).

**[Conclusion]** These results indicate that high-fat diets negatively affect bone health; however, these negative effects can be improved by exercises such as swimming.

**[Key words]** Swimming exercise, Bone mineral density, Microstructure, Trabecular bone, High-fat diet.

swimming exercise, and found that muscle contraction during swimming exercise can sufficiently improve bone mineral density<sup>14</sup>. However, with numerous suggestions that underwater exercises such as swimming, in which the load is reduced by buoyant force, hinder improvement in bone mineral density<sup>1, 24</sup>, further investigation is necessary to clarify the effects. Swimming exercise can be effective, as it not only improves metabolic disorders related to obesity by reducing the joint load and contributing to weight loss, but also positively affects bone health by improving bone mineral density.

The present study aimed to investigate the effects of swimming exercise for 8 weeks on bone mineral density and microstructural changes in the tibia and femur of rats.

## METHODS

### Animal models

Eight-week-old male Sprague–Dawley rats ( $n = 24$ ) were obtained from Damul Science Inc. (Daejeon, Korea). The rats were raised in a plastic cage (2 rats/cage) at the laboratory animal center of J University at 23–25°C and 70–80% humidity, with diurnal lighting.

### Methods

#### Dietary treatment

The animals were acclimated to the environment for 1 week. The rats were randomly assigned to a normal diet group ( $[n = 9]$  that was fed a diet containing 3.5% fat), and a high-fat diet group ( $[n = 15]$  that was fed a diet containing 45% fat). The rats were fed *ad libitum* for 8 weeks (Table 1). Three rats were sacrificed from each group to analyze the reduction in bone mineral density due to the high-fat diet. After 8 weeks of dietary treatment, rats were assigned to normal diet (NC,  $n = 6$ ), high-fat diet (HC,  $n = 6$ ), and high-fat diet + swimming exercise (HEX,  $n = 6$ ) groups. The experimental groups are described in <Fig. 1>.

#### Exercise method

Rats were subjected to low-intensity free swimming exercise without a load, similar in intensity to running

on a treadmill (13 m/min, grade 0%)<sup>32, 33</sup>. One week before the start of dietary treatment, rats underwent adaptation training (10–45 min/day) for 3 days in a water-filled (36°C ± 2°C) barrel (diameter 70 cm × height 70 cm). They were then subjected to swimming exercise (45–60 min/day, 5 times/week) for 8 weeks. The temperature of the water was maintained at 36°C ± 2°C, at which the core temperature is not affected<sup>14</sup>.

### Tissue collection

After 8 weeks of dietary treatment and exercise training, the rats were anesthetized with an intraperitoneal injection (1 mL/kg) of Zoletil (Virbac Laboratories, Paris, France), Rombun (Bayer Korea, Seoul, Korea), and saline solution mixed in a 2:1:2 ratio. Then, the tibias and femurs were separated and stored in a tube containing 4% formalin until the day of measurement<sup>28</sup>.


### Measurement parameters and method of analysis

#### Micro-computed tomography (CT)

To analyze tibial and femoral bone mineral density and microstructure, micro-CT (SkyScan 1076; Bruker, Kon-

**Table 1.** Composition of experimental diet

|                    | Normal diet | High-fat diet |
|--------------------|-------------|---------------|
| Casein             | 200.0       | 245.0         |
| L-Cystine          | 3.5         | 3.5           |
| Corn Starch        | 397.5       | 85.0          |
| Maltodextrin       | 35.0        | 115.0         |
| Sucrose            | 100.0       | 200.0         |
| Lard               | 20.0        | 195.0         |
| Soybean Oil        | 70.0        | 30.0          |
| Cellulose          | 50.0        | 58.0          |
| Mineral Mix        | 30.5        | 43.0          |
| Calcium Phosphate  | 3.5         | 3.4           |
| Vitamin Mix        | 19.0        | 19.0          |
| Choline Bitartrate | 2.5         | 3.0           |
| Red Food Color     | -           | 0.1           |
| Protein            | 20.5 %kcal  | 19.0 %kcal    |
| Carbohydrate       | 76.0 %kcal  | 36.2 %kcal    |
| Fat                | 3.5 %kcal   | 44.8 %kcal    |
| Total              | 100 %kcal   | 100 %kcal     |



tich, Belgium) was performed (voltage: 60kA, current: 167 $\mu$ A) through a 0.5-mm thick aluminum filter. The micro-CT images were reproduced in grayscale using Nercon Ver 1.3(SkyScan). The 2-dimensional (2D) images were reconstructed as 3D models by using CTAn and CTVox (SkyScan). A region of interest (ROI) between the growth plate of the femur or tibia to a point 0.5–4 mm away was analyzed using CTAn. Percent bone volume (BV/TV) of the ROI, which describes bone mineral density and microstructures, trabecular thickness (Tb.Th), trabecular number (Tb.N), and trabecular separation (Tb.Sp) were measured. On CTVox, images of 2D cross-sectional planes and 3D structures of trabecular bone within the femurs and tibias were created.

### Statistical analysis

The means and standard deviations of the collected data were calculated using SPSS Win 12.0. Weight differences between rats in the experimental groups after 8 weeks of a high-fat diet and swimming exercise were analyzed at different time points during the experiment using two-way repeated measures ANOVA. When interactions were observed between the main effects, one-way ANOVA was used to analyze the differences within and between groups at different time points. A t-test was used to determine the differences in bone mineral density between rats in the experimental groups after 8 weeks on a high-fat diet. One-way ANOVA was used to analyze the differences in the effects of swimming exercise after 8 weeks of a high-fat diet. The level of statistical significance was set at  $p < .05$ .

## RESULTS

### Weight changes due to high-fat diet and swimming exercise

Weight changes in the rats are shown in Table 2. Significant differences after 8 weeks of dietary treatment and swimming exercise were found at different time points within ( $F = 2901.837$ ) and between the groups ( $F = 186.486$ ) ( $p < .05$ ). A significant interaction was found between time point \* groups ( $F = 33.428$ ) ( $p < .05$ ). A post-hoc test, with which weight changes at different time points were compared, showed significant changes in the weight during the diet period and the final weight relative to the starting weight in rats in all three groups ( $F = \text{NC: } 3,704.059, \text{HC: } 6,755.808, \text{HEX: } 2,849.430$ ). Weight differences at different time points during the diet period were significantly higher in rats in the HC and HEx groups, which were assigned to a high-fat diet, than in those in the NC group, assigned to a normal diet ( $F = 106.378, p < .05$ ). The final weight of rats in the HC group was significantly higher ( $F = 49.684, p < .05$ ) than the final weight of rats in the NC and HEx groups. The final weights of rats in the HEx and NC groups were significantly lower than the final weights of rats in the HC group, indicating that 8 weeks of swimming exercise can decrease weight gained by a high-fat diet.

### Changes in bone mineral density and morphology due to a high-fat diet

Morphological changes in the tibias and femurs during 8 weeks of dietary treatment are shown in Fig 2.

**Table 2.** Change of body weight in normal diet control(NC), high-fat diet control(HC) and high-fat diet + swimming exercise(HEx) rats.

| Group    | Start weight(g) | Diet period weight(g) | Final weight(g) | F-value  |           |             |
|----------|-----------------|-----------------------|-----------------|----------|-----------|-------------|
|          |                 |                       |                 | Group    | Time      | Group* Time |
| NC(n=6)  | 162 $\pm$ 4.1   | 422 $\pm$ 4.8a        | 490 $\pm$ 9.5b  |          |           |             |
| HC(n=6)  | 160 $\pm$ 3.5   | 471 $\pm$ 5.1a#       | 547 $\pm$ 8.2b# | 186.485* | 2901.837* | 33.428*     |
| HEx(n=6) | 160 $\pm$ 4.4   | 474 $\pm$ 8.8a#       | 515 $\pm$ 11.1b |          |           |             |


Values expressed as mean $\pm$ standard error(g).

\*  $p < .05$

a  $p < .05$  vs. start weight in post hoc

b  $p < .05$  vs. diet period weight in post hoc

#  $p < .05$  vs. NC group in post hoc




**Figure 2.** Typical 2- and 3-dimensional images of femur and tibia in normal diet(A) and high-fat diet(B) feeding rats, as visualized using micro-computed tomography.


Significant morphological changes (reduced trabecular area and number) were observed in the 2D and 3D images of the tibias and femurs of rats from the high-fat diet group relative to those of rats in the normal group. Rats in the high-fat diet group had significantly lower bone mineral densities of the tibias ( $F = .645$ ) and femurs ( $F = .038$ ) than those of rats in the normal group ( $p < .05$ ) (Fig. 3).

### Changes in bone mineral density due to swimming exercise after a high-fat diet


Bone mineral densities of the femurs ( $F = 43.551$ ) and tibias ( $F = 20.843$ ) measured after 8 weeks of swimming exercise were significantly reduced in rats in the HC group relative to those of rats in the NC group ( $p < .05$ ). Similar results with regard to bone mineral density were observed between rats in the HEx group, which were subjected to swimming exercise, and those in the NC group (Fig. 5).



**Figure 3.** Micro-computed tomography derived trabecular bone mineral density at the femur and tibia in normal diet(ND) and high-fat diet(HFD) feeding rats. \* $p < .05$ .



**Figure 4.** Typical 2- and 3-dimensional images of femur and tibia in normal diet control(A), high-fat diet control(B) and high-fat diet + swimming exercise(C) rats, as visualized using micro-computed tomography.





**Figure 5.** Micro-computed tomography derived trabecular bone mineral density at the femur(A) and tibia(B) in normal diet control(NC), high-fat diet control(HC) and high-fat diet + swimming exercise(HEx) rats. \* $p < .05$ .

## Changes in trabecular microstructure after swimming exercise following a high-fat diet

Changes in trabecular microstructure in femurs and tibias after 8 weeks of swimming exercise are shown in Fig. 6 and Fig. 7, respectively. BV/TV ( $F = 8.216$ ) and Tb.N ( $F = 32.872$ ) of the femurs were significantly reduced in rats

in the HC group than in those in the NC group, significantly higher in rats in the HEx group than in the rats in the HC group, and similar to those of rats in the NC group, after 8 weeks of swimming exercise ( $p < .05$ ).

Tb.Th ( $F = 9.840$ ) and Tb.Sp ( $F = 8.110$ ) of femurs were significantly increased in rats in the HC group



compared to those in rats in the NC group, significantly decreased in rats in the HEx group compared to those in rats in the HC group, and similar to those of rats in the NC group, after 8 weeks of swimming exercise ( $p < .05$ ).

Significant differences in BV/TV ( $F = 9.383$ ) and Tb.N ( $F = 57.102$ ) of tibias were found between rats in the HC group and those in the NC and HEx groups. Significant differences of Tb.Th ( $F = 14.200$ ) and Tb.Sp ( $F = 23.733$ ) were found between rats in the HC group and those in the NC and HEx groups ( $p < .05$ ).

## DISCUSSION

Important factors associated with bone health and strength include not only bone mineral density but also characteristics of bone microstructure<sup>15, 16</sup>. Micro-CT is frequently used to assess bone microstructure, and is an effective method of measurement that requires lesser time than that required by histomorphometry; the use of micro-CT also prevents morphological changes in the sample during analysis<sup>31, 34</sup>. It provides structural information by enabling the quantification of bone area, bone volume fraction, and number, gap distance, and thickness of trabecular, as well as 3D reconstruction of bone microstructures<sup>9</sup>. Bone mineral density and structure of cancellous bone rather than cortical bone are known to determine bone strength. The proportion of cancellous bone and the number of trabecular are decreased throughout the bone area in patients with osteopenia or osteoporosis, while the gaps between trabecular are increased owing to the reduced microstructure in cancellous bone, with increased average trabecular thickness<sup>37</sup>. We aimed to analyze the effects of obesity and exercise on bone mineral density and microstructure of rats through micro-CT.

Recent studies on the correlation between obesity and bone health reported that bone mineral density decreases as the amount of body fat, blood cholesterol, neutral fat, and low-density lipoprotein cholesterol increase<sup>11</sup>, and as body mass index and waist circumference increase, leading to increased risk for osteoporosis<sup>25</sup>. In the present study, bone mineral density and the number of trabecular were significantly lower while trabecular thickness and gap distance were increased in rats in the high-fat diet group relative to those in rats in the normal diet group, indicating that weight gain negatively affects bone health (Fig. 2 & 3). Accumulation of adipose tissue as a result of obesity can disrupt the secretion of various substances involved in metabolism and inflammatory reactions<sup>3</sup>. An obesity-induced increase in the expression of regulators of inflammation such as interleukin (IL)-1, IL-6, and tumor necrosis factor- $\alpha$  induces osteoclast differentiation and bone resorption, and is cited as a cause of osteopenia and osteoporosis<sup>5, 8, 10</sup>. However, since an analysis of related factors (inflammatory cytokines, factors associated with bone resorption, etc.) was not performed in this study, it is difficult to identify

methods that can confirm changes in bone mineral density and an increase in the number of trabecular due to a high-fat diet. Further investigation is necessary in this regard.

Exercise training is commonly used to improve reduced bone mineral density, with weight training being the most common<sup>21, 22, 27, 36</sup>. Weight training increases mechanical load on the bones to stimulate bone metabolism and increase bone mineral density<sup>2</sup>. For obese patients, however, weight training can add excessive load on joints or bones, together with the increased body mass, and lead to joint injuries<sup>19, 35</sup>. Therefore, non-weight-bearing exercises such as swimming, water walking, and cycling that are easy on joints can be suggested as an effective exercise program for obese patients with reduced bone mineral density<sup>30</sup>. Non-weight-bearing exercises are commonly considered inappropriate for improving bone mineral density, since they do not sufficiently stimulate the bones. However, in a comparison of the effects of swimming and jumping in animal models with reduced bone mineral density (white mice) according to the degree of hindlimb suspension, bone mineral density increased after both types of exercises<sup>7</sup>, suggesting that differences according to the type of exercise may be insignificant. In addition, swimming exercise followed by resistance training without a load<sup>14</sup>, or with a load equivalent to 16% of body weight, can effectively increase the bone mineral density in the femur and the number of trabecular that had been reduced as a result of menopause. Therefore, differences in bone mineral density according to the intensity of swimming exercise may not be significant. In this study, 8-week swimming exercise without a load improved bone mineral density that was reduced owing to obesity, increased the bone area ratio and the number of trabecular, and decreased trabecular thickness and gap distance. This finding demonstrates that although the load on the bones is reduced owing to buoyant force during swimming exercise, stimuli produced through the continuous process of contraction and relaxation of skeletal muscle reach the bones and sufficiently stimulate bone metabolism<sup>14</sup>. However, mechanisms by which swimming exercise promotes bone metabolism have not yet been identified in this or other studies. Yang *et al.* reported a significant increase in blood irisin concentration after 8 weeks of swimming exercise without a load<sup>4</sup>, and increased expression of transcription factors involved in osteoclast differentiation in primary rat osteoblasts cells and MC3T3-E1 cells treated with irisin<sup>38</sup>. Further, administration of irisin in C57BL/6 mice for 4 weeks not only increases the expression of transcription factors related to osteoclast differentiation, but also increases cortical bone in the femur and tibia<sup>6</sup>. Therefore, an increase in the level of irisin after swimming exercise may be one of the potential stimulatory mechanisms of bone metabolism. However, the roles of swimming exercise and irisin in promotion of bone metabolism have not been clearly established in previous literature or in the

present study, which was aimed at analyzing the changes in bone mineral density and trabecular microstructure. Through additional research on the mechanism by which exercise training without a load improves bone mineral density, methods of improving negative effects on bones, such as reduced bone mineral density due to increased body mass, as well as safe and effective methods of exercise that can help obese patients maintain healthy bones may be established.

## ACKNOWLEDGMENTS

This research was supported by “Program of National University for Innovation and Transformation” funded by Chonbuk National University in 2015.

## REFERENCES

1. Abrahin O, Rodrigues RP, Marçal AC, Alves EA, Figueiredo RC, de Sousa EC. Swimming and cycling do not cause positive effects on bone mineral density: a systematic review. *Rev Bras Reumatol Engl Ed*. 2016; 56:345-351.
2. Aerssens J, Van Audekercke R, Geusens P, Schot LP, Osman AA, Dequeker J. Mechanical properties, bone mineral content, and bone composition (collagen, osteocalcin, IGF-I) of the rat femur: influence of ovariectomy and nandrolone decanoate (anabolic steroid) treatment. *Calcif Tissue Int*. 1993; 53:269-277.
3. Asayama K, Hayashibe H, Dobashi K, Uchida N, Nakane T, Koderia K, Shirahata A, Taniyama M. Decrease in serum adiponectin level due to obesity and visceral fat accumulation in children. *Obes Res*. 2003; 11: 1072-1079.
4. Yang XQ, Yuan H, Li J, Fan JJ, Jia SH, Kou XJ, Chen N. 2016. Swimming intervention mitigates HFD-induced obesity of rats through PGC-1 $\alpha$ -irisin pathway. *Eur Rev Med Pharmacol Sci*. 20(10):2123-2130.
5. Cao JJ. Effects of obesity on bone metabolism. *J Orthop Surg Res*. 2011; 15:6:30.
6. Colaianni G, Brunetti G, Faienza MF, Colucci S, Grano M. Osteoporosis and obesity: Role of Wnt pathway in human and murine models. *World J Orthop*. 2014; 5:242-246.
7. Falcai MJ, Zamarioli A, Okubo R, de Paula FJ, Volpon JB. The osteogenic effects of swimming, jumping, and vibration on the protection of bone quality from disuse bone loss. *Scand J Med Sci Sports*. 2015; 25:390-397.
8. Halade GV, Rahman MM, Williams PJ, Fernandes G. High fat diet-induced animal model of age-associated obesity and osteoporosis. *J Nutr Biochem*. 2010; 21:1162-1169.
9. Hanson NA, Bagi CM. Alternative approach to assessment of bone quality using micro-computed tomography. *Bone*. 2004; 35:326-333.
10. Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006; 444:860-867.
11. Hsu YH, Venners SA, Terwedow HA, Feng Y, Niu T, Li Z, Laird N, Brain JD, Cummings SR, Bouxsein ML, Rosen CJ, Xu X. Relation of body composition, fat mass, and serum lipids to osteoporotic fractures and bone mineral density in Chinese men and women. *Am J Clin Nutr*. 2006; 83:146-154.
12. Ip TY, Peterson J, Byrner R, Tou JC. Bone responses to body weight and moderate treadmill exercising in growing male obese (fa/fa) and lean Zucker rats. *J Musculoskelet Neuronal Interact*. 2009; 9:155-166.
13. Iwamoto J, Shimamura C, Takeda T, Abe H, Ichimura S, Sato Y, Toyama Y. Effects of treadmill exercise on bone mass, bone metabolism, and calcitropic hormones in young growing rats. *J Bone Miner Metab*. 2004; 22:26-31.
14. Ju YI, Sone T, Ohnaru K, Tanaka K, Fukunaga M. Effect of swimming exercise on three-dimensional trabecular bone microarchitecture in ovariectomized rats. *J Appl Physiol* (1985). 2015; 119:990-997.
15. Kleerekoper M, Villanueva AR, Stanciu J, Rao DS, Parfitt AM. The role of three-dimensional trabecular microstructure in the pathogenesis of vertebral compression fractures. *Calcif Tissue Int*. 1985; 37:594-597.
16. Koyama A, Kumasaka S, Kashima I. relationship between bone mineral density and 2D and 3D structure parameters of bone trabeculae. *Oral Radiol*. 2005; 21:62-68.
17. Kim M, Kim C, Choi YS, Kim M, Park C, Suh Y. Age-related alterations in mesenchymal stem cells related to shift in differentiation from osteogenic to adipogenic potential: Implication to age-associated bone diseases and defects. *Mech Ageing Dev*. 2012; 133:215-225.
18. Manolagas SC, Jilka RL. Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med*. 1995; 332:305-311.
19. Messier SP. Obesity and osteoarthritis: disease genesis and nonpharmacologic weight management. *Rheum Dis Clin North Am*. 2008; 34:713-729.
20. Muruganandan S, Roman AA, Sinal CJ. Adipocyte differentiation of bone marrow-derived mesenchymal stem cells: cross talk with the osteoblastogenic program. *Cell Mol Life Sci*. 2009; 66:236-253.
21. Nogueira JE, Branco LG, Issa JP. Bone repair: Effects of physical exercise and LPS systemic exposition. *Injury*. 2016; 47:1828-1834.
22. Notomi T, Lee SJ, Okimoto N, Okazaki Y, Takamoto T, Nakamura T, Suzuki M. Effects of resistance exercise training on mass, strength, and turnover of bone in growing rats. *Eur J Appl Physiol*. 2000; 82:268-274.
23. Novack DV, Teitelbaum SL. The osteoclast: friend or foe?. *Annu Rev Pathol*. 2008; 3:457-484.
24. Nyska M, Nyska A, Swissa-Sivan A, Samueloff S. Histomorphometry of long bone growth plate in swimming rats. *Int J Exp Pathol*. 1995; 76:241-245.
25. Park SJ, Ahn Y, Min HS, Oh KS, Park C, Cho NH, Kimm K. Osteoporosis prevalence of radius and tibia and related factors using multiple bone sites quantitative ultrasound measurement of the Korean health and genome study cohort women. *Korean J Community Nutr*. 2005; 10:536-545.
26. Robling AG, Castillo AB, Turner CH. Biomechanical and molecular regulation of bone remodeling. *Annu Rev Biomed Eng*. 2006; 8:455-498.
27. Rooney SI, Loro E, Sarver JJ, Peltz CD, Hast MW, Tseng WJ, Kuntz AF, Liu XS, Khurana TS, Soslowky LJ. Exercise proto-

- col induces muscle, tendon, and bone adaptations in the rat shoulder. *Muscles Ligaments Tendons J.* 2015; 4:413-419.
28. Ruan MZ, Dawson B, Jiang MM, Gannon F, Heggenes M, Lee BH. Quantitative imaging of murine osteoarthritic cartilage by phase-contrast micro-computed tomography. *Arthritis Rheum.* 2013;65:388-396.
  29. Sawant A, Chanda D, Isayeva T, Tsuladze G, Garvey WT, Ponnazhagan S. Noggin is novel inducer of mesenchymal stem cell adipogenesis: implications for bone health and obesity. *J Biol Chem.* 2012; 287:12241-12249.
  30. Shultz SP, D'Hondt E, Fink PW, Lenoir M, Hills AP. The effects of pediatric obesity on dynamic joint malalignment during gait. *Clin Biomech.* 2014; 29:835-838.
  31. Sone T, Tamada T, Jo Y, Miyoshi H, Fukunaga M. Analysis of three-dimensional microarchitecture and degree of mineralization in bone metastases from prostate cancer using synchrotron microcomputed tomography. *Bone.* 2004;35:432-438.
  32. Terada S, Goto M, Kato M, Kawanaka K, Shimokawa T, Tabata I. Effects of low-intensity prolonged exercise on PGC-1 mRNA expression in rat epitrochlearis muscle. *Biochem Biophys Res Commun.* 2002;16;296:350-354.
  33. Terada S, Tabata I. Effects of acute bouts of running and swimming exercise on PGC-1 $\alpha$  protein expression in rat epitrochlearis and soleus muscle. *Am J Physiol Endocrinol Metab.* 2004;286:E208-216.
  34. Thomsen JS, Ebbesen EN, Mosekilde L. Relationships between static histomorphometry and bone strength measurements in human iliac crest bone biopsies. *Bone.* 1998;22: 153-163.
  35. Williams F, Spector T. Recent advances in the genetics of osteoporosis. *J Musculoskelet Neuronal Interact.* 2006; 6:27-35.
  36. Wolff I, van Croonenborg JJ, Kemper HC, Kostense PJ, Twisk JW. The effect of exercise training programs on bone mass: a meta-analysis of published controlled trials in pre- and postmenopausal women. *Osteoporos Int.* 1999; 9:1-12.
  37. Won YY, Baek MH, Cui W, Kim MK. Analysis of the microstructural and mechanical properties in human femoral head trabecular bone with and without osteoporosis. *J Biomed Eng Res.* 2004; 25: 519-524.
  38. Qiao X, Nie Y, Ma Y, Chen Y, Cheng R, Yin W, Hu Y, Xu W, Xu L. Irisin promotes osteoblast proliferation and differentiation via activating the MAP kinase signaling pathways. *Sci Rep.* 2016; 6: 18732.